

Volume 9 Number 11 November 2018

# Indian Journal of Public Health Research & Development

An International Journal

#### SCOPUS IJPHRD CITATION SCORE

Indian Journal of Public Health Research and Development Scopus coverage years: from 2010 to 2017 Publisher: R.K. Sharma, Institute of Medico-Legal Publications ISSN:0976-0245E-ISSN: 0976-5506 Subject area: Medicine: Public Health, Environmental and Occupational Health CiteScore 2015-0.02
SJR 2015- 0.105
SNIP 2015- 0.034



Website: www.ijphrd.com



## Indian Journal of Public Health Research & Development

www.ijphrd.com

Contents	≣
Volume 9, Number 11 November 2018	8
1. The Spread Area of Malaria Vector in Timor Island, East Nusa Tenggara Province	. 1
Developing Community Resilience as a Supporting System in the Care of People with  Mental Health Problems in Indonesia	. 6
3. Effects of Knowledge of Vitamin D on Attitudes toward Sun Exposure among Middle-Aged and Elderly Indonesian Adults	11
4. The Effects of Age And Body Mass Index on Blood Glucose, Blood Cholesterol, and Blood Pressure in Adult Women	16
5. The Antioxidant Activity and Organoleptic Properties of Soursoup Leaf Tea (Annona Muricata L.) and Moringa Leaf (Moringa Oliefera L.) in Combination with Guava Leaf (Psidium Guajava)	22
6. Parent Communication Regarding Sexual and Reproductive Health of Adolescent:  A Qualitative Systematic Review	27
7. Five Types of Personality and the Locus of Internal Control in relation to Preeclampsia Pregnancy 3  Lusiana Meinawati, Kusnanto Kusnanto, Oedojo Soedirham	33
8. The Effect of Diabetes Self-Management Education, Based on The Health Belief Model, on the Psychosocial Outcome of Type 2 Diabetic Patients in Indonesia	37
9. How does the Dayak Ngaju Community Treat Malaria? A Qualitative Study on the Use of Traditional Medicine in Central Kalimantan Province, Indonesia	43
10. The Effect of Young Coconut Water against Morning Sickness among Women in the First Trimester of Pregnancy	48
11. Factors Affecting the Incidence of Chronic Energy Deficiency among Pregnant Women  Attending the Pulubala Community Health Centre	53
12. Implementation of Birth Preparedness and Complication Readiness (BPCR) in High Risk Pregnancies <i>Ika Mardiyanti, Nursalam, Arief Wibowo</i>	58

### Developing Community Resilience as a Supporting System in the Care of People with Mental Health Problems in Indonesia

#### Retno Lestari<sup>1</sup>, Ah Yusuf<sup>2</sup>

<sup>1</sup>Doctoral Program of Public Health, Faculty of Public Health, <sup>2</sup>Faculty of Nursing, Universitas Airlangga, Mulyorejo, Surabaya, Indonesia

#### **ABSTRACT**

People with mental disorders have complex disabilities which make them need other people's assistance in doing their daily living activities. Community resilience is one of the capacities which is needed to make a healthy environment for mental disorder patients. This study aims to describe how community resilience can be a supporting system in the care of people with mental disorders. A literature review was conducted in order to obtain 98 reference sources. Only 15 articles were eligible to be reviewed. The community resilience's supporting system consists of providing social, physical and economical infrastructures through employment opportunities or financial support and a decent life. Interactions between community members and people with mental disorders become a positive relationship, helping people with mental disorders to carry out their activities and to become independent. It also has an impact on decreasing the symptoms of relapse as well as reducing the level of dependency that cause them to continue to be a burden on their families, communities and global society.

Keywords: community resilience, mental health, supporting system

#### INTRODUCTION

The number of people with mental disorders, which is increasing with the years, is becoming a burden for families, communities and the global health care system. In 2016, the prevalence of psychiatric patients in the United States reached around 44.7 million people who were more than 18 years old. This number indicates that 18.3% of Americans suffer from a type of mental disorder. In Indonesia, the prevalence of severe mental disorders, such as schizophrenia, reached around 400,000 people, or as many as 1.7 per 1,000 population. In addition, the prevalence of emotional and mental disorders as indicated by symptoms of depression and anxiety for those aged 15 years old and over reached about 14 million people or 6% population.<sup>2</sup>

#### **Corresponding Author:**

Retno Lestari
Doctoral Program of Public Health,
Faculty of Public Health,
Universitas Airlangga, Mulyorejo, Surabaya, Indonesia
Email: retno.lestari-2017@fkm.unair.ac.id

The community is one of the support systems that can facilitate the recovery of mental disorders with the opportunity to live independently, to participate in community activities, to interact with others and to attach themselves to their daily environment. The community has an important role in creating positive support and wide opportunities for people with mental disorders for them to be fully involved in every event in the community.<sup>3–5</sup>

Previous studies explain that people with mental disorders also have awareness and the responsibility to deal with reality and to fulfil their actualisation. They understand expectations, goals, and responses to social needs and are able to work independently. They want to be recognised like any other individual who is able to adapt to society, despite experiencing inconsistencies in their way of thinking. If they only live in the hospital, what they see is only a group of people with mental disorders, and they can feel like life has stopped. Thus, support for people with mental disorders is needed, especially from the community around their homes. When they live with healthy individuals, they can plan for a better future.<sup>6,7</sup>

The treatment of mental disorders requires integration between adequate mental health services and

opportunities for the sufferers to develop themselves. The key points are the provision of information about their conditions, the opportunity to develop themselves, the chance to foster hope of them recovering from their mental disorders and the importance of mental health services' role in the community in providing positive emotional control for people with mental disorders.<sup>8,9</sup>

The existence of positive support from the community can prompt motivation in the patients for them to get up and recover from mental disorders. The forms of support obtained can be varied, such as the existence of positive assessments, being given the opportunity to tell stories, to be listened to by others and to able to engage in activities that exist in the community.7,10 recovery oriented experiences are required which recognise the holistic and diverse needs of individuals. Aim: The aim of the study was to examine the experiences of people living with a mental illness who participated in a recovery oriented program called Recovery Camp. The study aimed to examine how the program may have related and contributed to their mental health recovery. Methods: A descriptive phenomenological approach guided the study. Consenting participants (n = 5

The discovery of various factors affecting individuals and community mental health shows that people with mental disorders who live in the community can recover with community support. This study has intended to describe how community resilience can be used as a support system in the care of people with mental disorders.

#### **METHOD**

This study was a systematic review which used some of the literature from 98 articles obtained from Science Direct, Google Scholar, Proquest Health, Medical Complete, Proquest Nursing, Allied Health Source, Proquest Psychology Journals and Proquest Science Journals dating from 2008 to 2018. The inclusion criteria used were health or related research articles published in the last 10 years, research articles on topics about the community as a support system in the treatment of mental disorders, and research articles discussing the concept of community resilience.

Based on a literature review of 98 articles obtained from the listed databases, there were 50 articles that fulfilled the inclusion criteria. However, there were only 15 articles that discussed in detail about community resilience and the ability of the community to support the care of people with mental disorders.

#### RESULTS

Gillard et al. states that the care of mental illness sufferers has two main keys; a positive personal relationship (oneself) and wider social interactions (others). Self-awareness and positive self-acceptance give the individual the ability to reconcile themselves between the outside world and the internal mind. Self-recovery in mental disorders is the concept of balance between internal and external worlds. Table 1 shows the results of a literature review that discusses community resilience and community capacity in supporting the care of people with mental disorders.

#### DISCUSSION

Conflicts, stress and exposure to hazards can cause crises in the community system. Resilience is the ability of individuals or groups to react and move away from stress or exposure to stressors. Resilience describes the dynamic process of adapting to the crisis situation faced by the people. Resilience can be analogous to the philosophy of a bamboo plant, in which if it is exposed to the wind then it will bend and move, but in the end, it can return to its original position. 13

Community resilience is the method or ability of a community to show its resilience when facing existing stresses to restore their productivity through daily activities.<sup>13</sup> Resilient communities have become the nearest supporting system for people with mental disorders. Nowadays, resilience is defined as a process rather than a result. Resilience is the capacity of the system to face or recover from disturbances as well as changes that threaten adaptation or development functions. There are two stages in which resilience is considered to be a process, namely: 1) the disturbance or stressor that significantly affects the system and 2) the resilience capacity system that shows an ability to adjust and recover from the occurrence of trauma.<sup>14</sup>

The mechanism of resilience activation occurs throughout three stages, namely liminal suspension, compassionate witnessing and relational redundancy. Liminal suspension means the time of crisis in which individuals mutually reinforce each other in relation to their interpersonal relationships. Compassionate witnessing means how individuals within the

community are bound together and help to fulfil the needs of one another. Relational redundancy means coordination between the sources of the organisation concerned with strengthening resilience. When a critical situation arises, the group of people will gather, interact and perform in a variety of ways that can enable their existing capacity to survive and achieve resilience.<sup>15</sup>

Teo et al. found that the mechanism of resilience activation occurs during an exposure to stressor or crisis events. However, the existence of a stressor can generate resources in the community to strengthen interpersonal relationships, especially helping in the recovery of people with mental disorders in the community.<sup>16</sup>

Cohen et al. explained that community resilience involves local organisations and existing policies applied locally and regionally when strengthening resilience. Effective leadership is a leader who is able to interact with other community members and who can meet the needs desired by all members of the community. Community leaders have an important role in improving community resilience, primarily by focusing on the needs and demands of the local community.<sup>17</sup>

The involvement of a local government is needed in enhancing community resilience, including the development of access to technology that can support sources of social and economic capital. Increasing the productivity of community members through economics supported by local governments can increase resilience.<sup>18</sup>

There are several strategies for the reconstruction and revitalisation of sources of social capital after a crisis, including strengthening social networks and communities, building social organisations, and macro-social policies that can improve people's access to resources and power. In the aspect of social capital, social networks are important aspects that can affect mental well-being.<sup>3,19–21</sup>

Several previous studies have explained that economic resources are the most significant aspect in relation to building resilience. When individuals have suitable jobs and are in line with local needs, this will also increase the adaptive capacity of the community. A resilient community is a community that has the capacity to innovate and provide mutual support to other members of the community, including making a person with a mental disorder able to engage in productive activities in the community.<sup>10</sup>

Community resilience is an important focus of attention which can lead to positive attachments between people with mental and community disorders. Resilient communities are able to provide opportunities for sufferers to live like other individuals, who are able to be independent and work for themselves and others.

#### **CONCLUSIONS**

Community resilience is the capacity that is owned by the community that becomes an appropriate supporting system in the care of people with mental disorders living in the community. The increase in the provision of social, physical and economic infrastructure along with the interaction between community members and mental disorder patients is positive; helping people with mental disorders to carry out their daily activities and to become independent. This will also have an impact on decreasing the symptoms of relapse as well as reducing the level of dependency that will continue to be a burden on families, communities, and global society.

Conflict of Interest: None.

Ethical Clearance: Not required.

Source of Funding: Self-funded.

#### REFERENCES

- National Institute of Mental Health. Mental Illness [Internet]. National Institute of Mental Health. 2017 [cited 2017 Nov 29]. Available from: https://www.nimh.nih.gov/health/statistics/ mental-illness.shtml
- Kementerian Kesehatan Republik Indonesia.
   Riset Kesehatan Dasar. Jakarta, Indonesia; 2013.
- 3. Somasundaram D, Sivayokan S. Rebuilding community resilience in a post-war context: developing insight and recommendations a qualitative study in Northern Sri Lanka. Int J Ment Health Syst [Internet]. 2013 Jan 11 [cited 2018 Aug 15];7(1):3. Available from: http://ijmhs.biomedcentral.com/articles/10.1186/1752-4458-7-3
- 4. Piat M, Seida K, Sabetti J, Padgett D. (Em) placing recovery: Sites of health and wellness for individuals with serious mental illness in supported housing. Health Place [Internet]. 2017

- Sep [cited 2018 Aug 15];47:71–9. Available from: http://www.ncbi.nlm.nih.gov/pubmed/28759807
- Atallah DG, Contreras Painemal C, Albornoz L, Salgado F, Pilquil Lizama E. Engaging critical community resilience praxis: A qualitative study with Mapuche communities in Chile facing structural racism and disasters. J Community Psychol [Internet]. 2018 Jul 1 [cited 2018 Aug 15];46(5):575–97. Available from: http://doi. wiley.com/10.1002/jcop.21960
- 6. Mizuno E, Iwasaki M, Sakai I, Kamizawa N. Experiences of Community-Dwelling Persons Recovering From Severe Mental Illness. Arch Psychiatr Nurs [Internet]. 2015 Apr 1 [cited 2018 Aug 15];29(2):127–31. Available from: https://www.sciencedirect.com/science/article/pii/S0883941714001770
- 7. Picton C, Patterson C, Moxham L, Taylor EK, Perlman D, Brighton R, et al. Empowerment: The experience of Recovery Camp for people living with a mental illness. Collegian [Internet]. 2018 [cited 2018 Aug 15];25:113–8. Available from: https://doi.org/10.1016/j.colegn.2017.04.005
- 8. Bromley E, Gabrielian S, Brekke B, Pahwa R, Daly KA, Brekke JS, et al. Experiencing Community: Perspectives of Individuals Diagnosed as Having Serious Mental Illness. Psychiatr Serv [Internet]. 2013 Jul 1 [cited 2018 Aug 15];64(7):672–9. Available from: http://www.ncbi.nlm.nih.gov/pubmed/23545784
- Shepherd A, Sanders C, Shaw J. Seeking to understand lived experiences of personal recovery in personality disorder in community and forensic settings a qualitative methods investigation. BMC Psychiatry [Internet]. 2017 [cited 2018 Aug 15];17(1):282. Available from: http://www.ncbi.nlm.nih.gov/pubmed/28764672
- Hills JM, Michalena E, Chalvatzis KJ. Innovative technology in the Pacific: Building resilience for vulnerable communities. Technol Forecast Soc Change [Internet]. 2018 Apr [cited 2018 Aug 15];129:16–26. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0040162517302093
- 11. Gillard S, Turner K, Neffgen M. Understanding recovery in the context of lived experience of personality disorders: a collaborative, qualitative research study. BMC Psychiatry [Internet].

- 2015 Dec 31 [cited 2018 Aug 15];15(1):183. Available from: http://www.ncbi.nlm.nih.gov/pubmed/26227023
- 12. Henry CS, Sheffield Morris A, Harrist AW. Family Resilience: Moving into the Third Wave. Fam Relat [Internet]. 2015 Feb 1 [cited 2018 Aug 15];64(1):22–43. Available from: http://doi.wiley.com/10.1111/fare.12106
- 13. Kirmayer LJ, Sehdev M, Whitley R, Dandeneau SF, Isaac C. Community Resilience: Models, Metaphors and Measures. Int J Indig Heal [Internet]. 2009;5(1):62–117. Available from: http://journals.uvic.ca/index.php/ijih/article/view/12330
- 14. Abramson DM, Grattan LM, Mayer B, Colten CE, Arosemena FA, Bedimo-Rung A, et al. The resilience activation framework: a conceptual model of how access to social resources promotes adaptation and rapid recovery in post-disaster settings. J Behav Health Serv Res [Internet]. 2015 Jan [cited 2018 Aug 15];42(1):42–57. Available from: http://www.ncbi.nlm.nih.gov/pubmed/24870399
- 15. Powley EH. Reclaiming resilience and safety: Resilience activation in the critical period of crisis. Turner N, Gray GC, editors. Hum Relations [Internet]. 2009 Sep 13 [cited 2018 Aug 15];62(9):1289–326. Available from: http://journals.sagepub.com/doi/10.1177/0018726709334881
- 16. Teo WL, Lee M, Lim W-S. The relational activation of resilience model: How leadership activates resilience in an organizational crisis. J Contingencies Cris Manag [Internet]. 2017 Sep 1 [cited 2018 Aug 15];25(3):136–47. Available from: http://doi.wiley.com/10.1111/1468-5973.12179
- 17. Cohen O, Goldberg A, Lahad M, Aharonson-Daniel L. Building resilience: The relationship between information provided by municipal authorities during emergency situations and community resilience. Technol Forecast Soc Change [Internet]. 2017 Aug 1 [cited 2018 Aug 15];121:119–25. Available from: https://www.sciencedirect.com/science/article/pii/S0040162516306564

- 18. United Nations Office for Disaster Risk Reduction. Sendai Framework for Disaster Risk Reduction 2015 - 2030 [Internet]. Geneva; 2015 [cited 2018 Aug 15]. Available from: https://www.preventionweb.net/files/43291\_ sendaiframeworkfordrren.pdf
- 19. McAneney H, Tully MA, Hunter RF, Kouvonen A, Veal P, Stevenson M, et al. Individual factors and perceived community characteristics in relation to mental health and mental well-being. BMC Public Health [Internet]. 2015 Dec 12 [cited 2018 Aug 15];15:1237. Available from: http://www.ncbi.nlm.nih.gov/pubmed/26655388
- 20. Patel RB, Gleason KM. The association between social cohesion and community resilience in two urban slums of Port au Prince, Haiti. Int J Disaster Risk Reduct [Internet]. 2018 Mar 1 [cited 2018 Aug 15];27:161–7. Available from: https://www.sciencedirect.com/science/article/ pii/S2212420917302947
- 21. Pfefferbaum RL, Pfefferbaum B, Zhao YD, Van Horn RL, McCarter GSM, Leonard MB. Assessing community resilience: A CART survey application in an impoverished urban community. Disaster Heal [Internet]. 2016 [cited 2018 Aug 15];3(2):45–56. Available from: http://www.ncbi.nlm.nih.gov/pubmed/28229014